

**Lin v. MetLife**

**07 civ. 3218**

## **EXHIBIT K**

Metropolitan Life Insurance Company  
One MetLife Plaza  
27-01 Queens Plaza North  
Long Island City, New York, NY 11101

**MetLife®**

**Tomasita L. Sherer**

Senior Counsel  
Law Department  
[tsherer@metlife.com](mailto:tsherer@metlife.com)  
Tel 212 578-3102 Fax 212 743-0668

May 16, 2008

**VIA FEDERAL EXPRESS**

Eric Dinnocenzo, Esq.  
TRIEF & OLK  
150 E. 58<sup>th</sup> Street, 34<sup>th</sup> Fl.  
New York, NY 10155  
(212) 486-6060 (Tel)


**Re: Lin v. MetLife, Index No. 07-CV-3218 (S.D.N.Y.)**  
**Policy No. 204 126 416 ET**

Dear Eric:

Enclosed please find company guidelines responsive to your requests, bates stamped  
CONFIDENTIAL ML 1091-1106. Please utilize in accordance with our confidentiality  
agreement. Thank you.

If you have any questions, please feel free to call me.

Sincerely yours,



Tomasita L. Sherer

Enclosures

## Cholesterol (Hyperlipidemia, Hypercholesterolemia)

Text

### Lipids

Total Cholesterol mg/dl	Cholesterol / HDL Ratio			
	< 3.5	3.5 - 6.0	6.1 - 10	> 10
< 200	0	0	0	Refer to Med Dir
200 - 240	0	0	0	+25
241 - 300	0	0	0	+50
301 - 350	0	0	+50	+100
351 - 400	0	+50	+100	+200
> 400	Refer to Med Dir	Refer to Med Dir	Refer to Med Dir	Refer to Med Dir

Cholesterol HDL Ratio not  
available

### Isolated Triglyceride Elevations

Triglycerides may be elevated in a nonfasting specimen. Fasting status should be taken into account when evaluating isolated triglyceride elevations.

≤400	+0
401 - 800	+50
> 800	Refer to Med Dir

(VERSION NOTE This topic replaces and supersedes its previous version OUTDATED-Cholesterol Rating Schedule V0 as of April 3, 2001)

## Liver Function Tests

**Other Tests**

**Medications**

**Info**

**Text**

### Key Factors

**Alkaline phosphatase** Normal range 30 - 115 U/L

No other LFT abnormalities or history of malignancy

#### Adolescents

To 345 (or 3 times lab normal)

+ 0 (May be considered for Select Preferred/Elite)

346 up

Refer to Med Dir

#### Pregnant Females

To 230 (or 2 times lab normal)

+ 0 (May be considered for Select Preferred/Elite)

231 up

Refer to Med Dir

#### All Others

During healing phase of bone fracture or following bone surgery

I.C.

Otherwise:

To 185

+ 0

186 - 230

+ 50 to + 100

231 up -

Refer to Med Dir (Usually Decline)

With other LFT abnormalities or history of malignancy

May require current evaluation

Refer to Med Dir

Sum ratings to Decline

**Bilirubin** Normal range 0.5 - 1.5

Cause unknown

Up to 3.5 mg/dl

+ 0

Over 3.5 mg/dl - Refer to Med Dir

+ 50 to 100 up

Cause known

All

Rate for cause

### GGT/ ALT/ AST Ratings

(VERSION NOTE This topic replaces and supersedes its previous version OUTDATED-Liver Function Tests Rating Schedule V0 as of February 8, 2003)